

EFFECTS OF AGRO-CLIMATIC VARIABLES ON YIELD OF SELECTED VARIETIES OF RICE IN TARABA STATE, NIGERIA

A. S. Umar*¹, B. A. Ankidawa², A. A. Adebayo¹ and A. Ezra¹

¹Department of Geography, School of Environmental Sciences, Modibbo Adama University of Technology, Yola

²Department of Agric. and Environmental Engineering, School of Engineering and Engineering Technology, Modibbo Adama University of Technology, Yola

*Correspondent author: sadiqumar2001@yahoo.com

ABSTRACT

Rice is one of the major crops cultivated in Nigeria and production of the crop is being affected by many factors among which are climatic parameters. The aim of this study is to examine the effect of climatic variables on rice yield in Taraba State. Agro-Climatic variables and rice yield (Local, Ofada, FARO 44 and FARO 58 varieties) were the data used. Stepwise regression was used to analyze the data. Result on the effect of climatic variables on local rice variety revealed that, mean annual temperature was identified to be the critical climatic factor affecting the yield of the variety and it is statistically significant at $p < 0.01$, and accounted for 43.1% to the variation in Local Rice yield in the State. Minimum Temperature, Solar Radiation and June Rainfall were identified to be the critical climatic factors affecting the yield of Ofada rice variety and the variables were statistically significant at $p < 0.01$ and accounted for 64.0% to the variation in yield of Ofada rice variety in the state. Prediction model for FARO 44 rice reveals that, Mean temperature, August and September rainfall were identified to be the critical climatic factors affecting the yield of the variety in the State and the variables were highly significant at $p < 0.01$ and accounted for 61% to the variance in the yield of the variety. In the case of FARO 58, it was observed that, mean temperature, August and October rainfall were identified to be the critical climatic variables affecting the yield of the variety in the state and were highly significant at $p < 0.01$ and accounted for 71.2% to the variation in yield of FARO 58. based on the findings of this studies, it was recommended that, cultivation of local, FARO 44 and 58 rice varieties should be encouraging in places with high temperature and August rainfall amount and low September and August rainfall in the state. Good and better adaptation method should be practice by the farmers to help against unforeseen event that will affect their crop production. In addition, there is a need for government assistance in terms of improved varieties of rice and other input. Use of information technology by farmers will greatly facilitate communication among farmers on new method of farming.
Key words: Effects, Agro-Climatic variables, Rice yield, FARO 44, FARO 58, Taraba State

1 INTRODUCTION

Rice is one of the major crop cultivated in Nigeria, the crop plays a very significant role in food security and provision of employment but its production is faced with some challenges among which is climate. Climate is very sensitive to all stages of crop production; from land clearing and preparation, through crop growth and management to harvesting, storage, transportation and marketing of the product [1]. The amount and distribution of Precipitation, solar radiation, wind, temperature, relative humidity and other climatic parameters affect and solely influence the global distribution of crops as well as their productivity and also determine the types of crop a farmer cultivate, the yield of the crop and hence farmers' profit [1, 2, 3, 4]. In addition, the amount of Sunlight, temperature, moisture and carbon dioxide (CO₂) determines seed germination, the time emergence of plant, the rate of growth of roots, stems, and leaves and flower development. They also determine when plant produces flowers, and consequently the filling of grain or the expansion of fruits [5]. In Taraba State, the Government has designed programs on food security in the past to meet up with the demands of the growing population. These programs include; the integrated rural development program

which is anchored on the Agricultural Development Program (ADP), Agricultural Mechanization Program in 2005, Taraba State palm oil production program in 2001, Taraba State fertilizer voucher system policy in 2010, Taraba State policy on tractorization in 2007 and Taraba State agricultural produce and marketing agency TAPMA 2007 which focused on maize and rice production [6]. Although these programs were set to boost agricultural practices in the state, but farmers are still battling with some challenges among which are climate related issues in their farming operation [7, 8, 9]. In respect to this, there is a need to examine the critical climatic variables affecting different varieties of rice in the State.

2 STUDY AREA

Taraba State was carved out of the former Gongola State on 27th August 1991 by the then regime of General Ibrahim Babangida. The State is one of the Nigerian thirty-six (36) state located in North-Eastern part of the country and has a coordinate of latitude 6°30' and 8°30' North of equator and longitude 9°00' and 12°00' East of the Greenwich meridian (Figure 1). The area is made up of high plains which covered those parts of the Benue low lands lying above flood level but below 1000-foot contour line and include places around Karim Lamido, Jalingo, Sunkani and some part of Wukari while the high highlands are erosional in nature and are cut in sedimentary formations. River Benue is the major river in the state [10]. River Donga and Taraba are the dominant river systems which flow across the Muri plains to drain the entire State together with the minor ones, such as the Lamorde and Mayo Ranewo [11].




Figure 1. Map of Tatraba State showing the study area

Climate of the State is mainly influenced the rain-bearing south-west air mass and the dry dusty north-east trades or harmattan. Rainy season in Mambilla Plateau lasts from February to November with a mean annual rainfall of about 1,850 m, while at the other part of the state lasted from April to October with mean annual rainfall varying between 1058mm around Jalingo and Zing, to about 1300 mm around Serti and Takum [12]. Temperature during rainy season in Mambilla drops to as low as 15°C while the mean annual temperature around Jalingo is about 28°C with maximum temperature varying between 30°C and 39.4°C and minimum temperatures range between 15°C to 23°C [12]. Alluvial soil types are found on the flooded plains of rivers they run along Benue River and other rivers, and do not depend highly on climate and vegetation for their formation but their underlying parent rock is the most important factor in their formation [13]. Sudan Vegetation, Northern Guinea Savanna, Southern Guinea Savanna, Forest derive savanna and mountain forest and grassland are the major vegetation types in the area. Sudan vegetation covered places around Karim Lamido, Lau, Jalingo, Ardo Kola, Yorro and Zing LGA, while Northern and southern savanna covers the major part of the State and include LGAs such as Gassol, Ibi, Wukari, Donga, Bali, Takum, Ussa, Kurmi and Gashaka LGA.

3 MATERIALS AND METHOD

Annual rainfall, monthly rainfall, dry spell, onset date of rain, cessation date of rain, Length of Rainy Season (LRS), seasonality index and hydrological ratio were extracted from the daily, monthly and annual rainfall for a period of 38 years (1979 to 2017) while, monthly and annual minimum temperature, maximum temperature, mean temperature, relative humidity, solar radiation and wind speed for the period 1979 to 2015 which is the longest period for which records of the variables are available in the State were obtained from Upper Benue River Basin Development Authority (UBRBDA), Taraba State Agricultural Development Program (TADP) (Area, Zonal and Head office) in the State, Taraba State University Jalingo, Federal Polytechnic Bali and Nigerian Meteorology (NIMET) Ibi. Rice yield data based on varieties (Local, Ofada, FARO 44 and FARO 58 varieties) and LGAs on the other hand were collected from Taraba State Agricultural Development Program (TADP) Area, Zonal and Head office in the State for the available years (2011-2017). The varieties were selected based on the fact that they are the major varieties of rice cultivated in all the Local Government Areas of the State. The yield records were collected by the staff of TADP annually through a survey method called Crop Area Yield Survey (CAYS). In this survey method, rice yield record based on varieties and cultivated area of rice farmers in each LGA of the State were collected, analyzed, summarized and recorded in tons per hectare. According to [1], there are two methods of crop-climate study; Simulation or physiological method and statistical or empirical method. Simulation or physiological method examines the impact of climatic factors on physiological characteristics of plant (Number of leaves, leave length, stem diameter, plant height among others) while statistical or empirical method studies the relationship between climatic factors and crop yield over time. In regards to the aim of this research and the data required to achieve this objective, the second method was used to examine the impact of climatic variables on rice yield which was also used by many authors in their analysis [1, 14, 15, 16, 17, 18, 19]. In other to examine the effect of climatic variables on rice yield over time and space, Station-Year method suggested by [15] as the best and accurate method of studying spatio-temporal impact of agro-climatic variables on rice yield was used. Applying the method, Agro-climatic variables (minimum, maximum and mean temperature, relative humidity, solar radiation, wind speed, dry spell, onset, cessation, LRS, seasonality index, hydrological ratio among others) as well as rice yield were arranged based on the available stations (LGAs) and years of record in the state for efficient regression analysis. This method was conducted on all the selected rice varieties using SPSS and MiniTab statistical packages. Regression analysis was used to examine the relationship between the climatic variables and yield of different varieties of rice as well as the effect of the climatic variables on rice yield. The Statistical Regression model used in this research is presented below;

$$Y = a + b_1X_1 + b_2X_2 + b_3X_3 \quad (1)$$

Where: Y = Rice yield, X_1 = Mean Temperature (For Local Rice), X_1 = Minimum Temperature, X_2 = Solar Radiation and X_3 = June Rain (for Ofada), X_1 = Mean Temperature, X_2 = August Rain and X_3 = September Rain (for FARO 44), X_1 = Mean Temperature, X_2 = August Rain and X_3 = October Rain (for FARO 58), a and b are constant.

4 RESULTS AND DISCUSSION

4.1 Effect of Agro-climatic Variables on Local Rice Variety

Table 1 showed the effect of critical agro-climatic elements on Local Rice yield in the study area. The result revealed that the Mean temperature is the only agro-climatic variable that is highly significant to the cultivation of Local rice yield in the State. The prediction model showed a significant F-value of 28.726 at $p < 0.01$ and accounted for 43.1% of the variation in Local Rice yield in the State. Mean Temperature as the only critical climatic element that is highly significant to Local rice yield showed a positive significant T-value of 5.36 at $p < 0.01$. The coefficient value of the critical element is 0.526 which implies that a unit increase in Mean Temperature will lead to an increase in variation of Local Rice yield of 0.526 in the State while holding other variables constant. Prediction of mean temperature as the critical climatic element influencing Local rice yield in the State also suggested that Local rice cultivation is highly determined by the sufficiency of Mean temperature for its effective production, and implies that places of high mean temperature in the state are good for cultivation of local rice variety while places with low mean temperature amount are detrimental to local rice cultivation.

in the state. This is not surprising because Mean Temperature is one of the critical climatic elements that influence rice yield at the different developmental stage of its production and the optimum temperature for normal development of rice ranges from 27°C to 32°C [20, 21, 22]. This result also agrees with the findings of [16].

Table 1. Stepwise Regression result between Local Variety and Agro-Climatic Variables

Predictor	Coef	SE Coef	T	R ² (%)	R ² (adj)	R ² (pred)	VIF
Constant	-11.971	2.671	-4.48**				
Mean Temp	0.52644	0.09816	5.36**	43.1	41.6	36.90	1.000

** T-value is significant at 1%; Local Variety = - 12.0 + 0.526 Mean Temperature

4.2 Effect of Agro-Climatic Variables on Ofada Variety

Table 2 showed the critical agro-climatic elements that influence the yield of Ofada rice variety. All the selected Agro-climatic variables were subjected to stepwise multiple regressions and the model selected three (June Rainfall, Minimum Temperature and Solar Radiation) climatic variables that are highly significant to Ofada rice yield in the State. Selection of June Rainfall, Minimum Temperature, and Solar Radiation showed that the variables are the critical contributors to variation in yield of Ofada rice variety in the state and their variation over time and space influences the growth and yield of Ofada rice in the State. The model is statistically significant with F-value = 14.21 at p<0.01 and accounted for 64.0% of the variance in yield of Ofada rice variety in the state (R²= 0.6398, Adjusted R² = 0.5945). Solar Radiation has the strongest negative contribution of 0.7836 to variation in yield of Ofada rice followed by Minimum Temperature with a positive contribution of 0.3384 and then June Rainfall with a negative contribution of 0.0033. The positive contribution of Minimum Temperature to the yield of Ofada rice variety suggested that a unit increase in Minimum temperature will lead to an increase in yield of Ofada rice with 0.3384, while a decrease in its amount will decrease the yield of the variety. Unlike the Minimum Temperature, June rainfall and Solar Radiation showed a negative contribution of -0.0033 and -0.7836 respectively, this negative contribution suggested that a unit increase in June rainfall and Solar Radiation will decrease the yield of Ofada rice yield with 0.0033 and 0.7836 respectively. In addition, the 64% contribution of the Agro-climatic variables to yield of Ofada rice signifies that growth and yield of Ofada rice is strongly determined by the amount of these three parameters and place of high minimum temperature will increase Ofada rice yield while places of high Solar Radiation and June Rainfall will reduce the yield of Ofada rice variety in the State. This is not surprising because heat parameters like Minimum temperature and solar radiation play a significant role in the growth and yield of rice [23]. Moreover, the June rainfall coincides with the planting to seedling (Nursery) stage of rice development in the State, as such, places with excess rainfall will make the rice plot to be submerged with water thereby making the place to be detriment to Ofada rice development since water requirement for rice production at planting to seedling is low [24, 25]. In addition, excess rainfall at planting to the emergence stage of rice will make the rice field to be submerged for a long period of time which leads to the deficiency of essential nutrients like nitrogen, magnesium, potassium, and calcium required for plant growth [26].

Table 2. Stepwise Regression result between Ofada Variety and Agro-Climatic Variables

Predictor	Coef	SE Coef	T	R ² (%)	R ² (adj)	R ² (pred)	VIF
Constant	13.119	6.171	2.13*				
Min. Temp	0.33836	0.09385	3.61**	40.70	38.42	33.31	1.009
Solar Rad.	-0.7836	0.2362	-3.32**	55.17	51.58	47.48	1.173
June Rain.	-0.0033	0.00137	-2.42*	63.98	59.45	53.06	1.164

** T-value is significant at 1%, * T-value is significant at 5%

Ofada = 13.1 + 0.338 Minimum Temperature- 0.784 Solar Radiation - 0.00333 June Rainfall

4.3 Effect of Agro-Climatic Variables on FARO 44 Variety

Table 3 displayed the stepwise regression result between the agro-climatic variables and FARO 44 rice yield. The prediction model contains three predictors and showed a significant F-value of 19.117 at $p < 0.01$ and accounted for 61.4% of the variance in the yield of FARO 44 rice in the State. The model also showed a significant positive T-value of 5.981 and 3.440 at $p < 0.01$ for Mean Temperature and August rainfall respectively, while September rainfall showed a significant negative T-value of 2.592 at $p < 0.05$. This result implies that Mean Temperature contributed high to variation in yield of FARO 44 in the State with a coefficient value of 0.5013 and R^2 to 46.44%, followed by August rainfall with a coefficient of 0.0044 and the R^2 increases from 46.44% to 54.26%. September rainfall is the only parameter that showed a negative coefficient of 0.0026 and increased the R^2 to 61.44%. This climatic contribution explained that a unit increase in Mean Temperature and August rainfall will lead to an increase in yield of FARO 44 rice of 0.5013 and 0.0044 respectively, while a unit increase in September rainfall will reduce the yield of the variety with 0.0033. In addition, the result clearly suggested that places with high mean temperature and August rainfall support the growth and yield of FARO 44 but places with a low amount of the parameters are detriment to the growth and yield of the rice variety. This is not surprising because places with low mean temperature amounts of $< 20^\circ\text{C}$ are not suitable for the growth and yield of rice [27]. The negative contribution of September rainfall, on the other hand, suggested that places of high September rainfall are detriment for growth and yield of FARO 44 while places with low September Rainfall are favorable for FARO 44 cultivation. This relationship is not coincident because FARO 44 (SIPI) has a maturity period of 90 to 100 days and given that rice planting in the State started from the month of June, this clearly suggest that September is the maturity period of FARO 44 and maturity period of rice required less or no rainfall especially in the last 15 to 20 days where water from the farm plot is drained completely. In addition, rice variety mature within stipulated time frame and any supply of climatic amount after it maturity will be insignificant to it yield since rice growth is very sensitive to climate at different growth stages [20, 28, 29]. The positive contribution of August rainfall is also not surprising because the month of August corresponds with the critical development stage of rice in the study area (Reproductive Stage) where rainfall amount is very critical to rice growth. The stage is recognized to be critical because a high rainfall amount in the stage determines the effectiveness of the stage and any unfavorable rainfall supply will damage the crop yield [24, 25].

Table 3. Stepwise Regression result between Faro 44 Variety and Agro-Climatic Variables

Predictor	Coef	SE Coef	T	R^2 (%)	R^2 (adj)	R^2 (pred)	VIF
Constant	-11.508	2.312	-4.98**				
Mean temp	0.50128	0.08379	5.98**	46.44	45.03	40.39	1.027
August Rain	0.0044	0.001285	3.44**	54.26	51.79	42.85	1.143
Sept. Rain	-0.0026	0.0009975	-2.59*	61.44	58.23	46.98	1.163

** T-value is significant at 1%, * T-value is significant at 5%

FARO 44 = - 11.5+ 0.501 Mean Temperature+ 0.00442 August Rain - 0.00258 September Rain

4.4 Effect of Agro-Climatic Variables on FARO 58 Variety

Table 4 showed the stepwise regression result between FARO 58 and the agro-climatic variables in the state. The result explained that the prediction model contains three different climatic variables that are critical in determining the variation in yield of FARO 58 rice variety. The model presents the percentage contribution of the selected climatic variables to variation in yield of FARO 58 variety. It also predicts the yield of FARO 58 as a result of a unit increase in the selected climatic variables in the study area. The model revealed that the Mean temperature, August Rainfall, and October Rainfall are the highly significant climatic variables that determine the variation in yield of FARO 58 rice variety in the study area. The model predicted that the three variables are highly significant with an F-value of 21.42 at $p < 0.01$ and accounted for 71.2% to the variation in yield of FARO 58 in the study area. The model also revealed that the Mean temperature and August rainfall contributed positively to the yield of FARO 58 while October rainfall contributed negatively to the yield of the crop. The positive

contribution of Mean Temperature and August Rainfall suggested that a unit increase in mean temperature and August rainfall will lead to an increase in yield of FARO 58 rice variety with 0.351 and 0.008 respectively while the negative coefficient of October rainfall implies that a unit increase in rainfall amount in the month of October will decrease the yield of FARO 58 with 0.0039 in the State. In addition, the positive effect of Mean Temperature and August Rainfall suggested that places of high temperature and August rainfall are favorable to the cultivation of FARO 58, while places of high rainfall amount in the month of October is not favorable to FARO 58 rice cultivation. Rainfall amount in the month of August coincides with the vegetative stage of rice growth in the State where water requirement at this stage is high compare to other stages and any deficiency in rainfall amount within the stage will damage rice yield [24, 25]. Unlike the month of August, the month of October is marked as the end of the growing season and harvesting period in the State. This stage is recognized to be the maturity stage of rice growth where a high rainfall amount is a detriment to rice. In fact, during this stage especially in the last 15 to 20 days before harvesting, water is not required; instead, all water in the farm is drained out.

Table 4. Stepwise Regression result between Faro 58 Variety and Agro-Climatic Variables

Predictor	Coef	SE Coef	T	R ² (%)	R ² (adj)	R ² (pred)	VIF
Constant	-8.029	2.972	-2.70*				
Mean Temp	0.3513	0.1044	3.36**	38.56	36.37	30.53	1.306
August Rain	0.007772	0.001702	4.57**	63.89	61.22	56.23	1.304
October Rain	-0.0039	0.001524	-2.57*	71.19	67.87	64.63	1.027

** T-value is significant at 1%, * T-value is significant at 5%

FARO 58 = - 8.03 + 0.351 Mean Temperature + 0.00777 August - 0.00391 October

4.5 Model Validation

The results of model validation shows that, mean temperature is one of the critical agro-climatic elements that positively influence the yield of Local, FARO 44 and 58 rice varieties in the State as shown in Table 5. The nature of the relationship clearly suggested that places with high mean Temperature amount in the State support the growth and yield of rice compared to places with low mean temperature which is a detriment to rice yield in the state. This is not surprising because temperature is identified to be one of the critical climatic elements that influence growth and yield of rice crop at different growth stage of rice and the optimum temperature requirement is between 27°C to 32°C [17, 18, 29]. Similarly, rainfall amounts in the months of June, August, September, and October during the growing period of the crop are critical and determine the yield of the selected varieties of rice in the State. Following the result of the analysis obtained, it is clear that annual rainfall is not a major determinant of the growth and yield of the varieties in the State. This is not surprising because it is not the volume of rainfall received in a year that determined the yield of rice, but how efficient is the rainfall amount distributed at the various developmental stages of the crop especially the critical stages of the crop development. In the same vein, rice yield is not determined by the volume of rainfall received in a year but the efficiency of the rainfall at different developmental stages of the crop growth [30, 31, 32, 33, 34].

Table 5. Summary of the Identified Critical Agro-Climatic Variables based on Rice Varieties

Rice varieties	Critical Climatic Variables	Nature of the Relationship
Local Variety	i. Mean Temperature	Positive
	i. Minimum Temperature	Positive
Ofada	ii. Solar Radiation	Negative
	iii. June Rainfall	Negative
	i. Mean Temperature	Positive
FARO 44	ii. August Rainfall	Positive
	iii. September Rainfall	Negative
	i. Mean Temperature	Positive
FARO 58	ii. August Rainfall	Positive
	iii. October Rainfall	Negative

5 CONCLUSION

Following the result on the critical agro-climatic variables influencing rice yield in the State, it was also concluded that Mean annual temperature is critical to growth and yield of local rice variety in the State, while, Minimum temperature, June rainfall and Solar Radiation are critical to growth and yield of Ofada rice variety. Mean annual temperature, total August rainfall and September rainfall on the other hand are the major climatic parameters that are critical to growth and yield of FARO 44, while FARO 58 growth and yield are majorly determined by Mean annual temperature, total August and October rainfall. It was also concluded that, annual rainfall is not a major determinant of the growth and yield of the varieties in the State. This is based on the fact that, it is not the volume of rainfall received in a year that determines the yield of rice, but how efficient is the rainfall amount distributed at the various developmental stages of the crop especially the critical stages of the crop development.

REFERENCE

1. Ayoade, J.O. (2005). Introduction to Agroclimatology. *Vantage, Publisher, Ibadan, Nigeria*, pp. 113-117.
2. Stigter, C.J. (2004). The Establishment of needs for Climate Forecasts and other Agrometeorological Information for Agriculture by Local, National and Regional Decision-Makers and Users Communities. In: *Applications of Climate Forecasts for Agriculture. AGM-7/WCAC-1, WMO/TD-No. 1223, WMO, Geneva*, pp. 73–86.
3. Ankidawa, B.A., Tanko, J.A. and Umar, A.S. (2015). Climatic Variability and River Benue Discharges in Yola and Garoua, West Africa. *Journal of Environment, Ecology, Family and Urban Studies*, vol. 5, Issue 1, pp. 1-14.
4. Umar, A.S. and Ankidawa, B.A. (2016). Climate Variability and Basin Management: A Threat to and from Wetlands of Komadugu Yobe Basin, North Eastern Nigeria. *Asian Journal of Engineering and Technology*, vol. 4, Issue 2, pp. 25-36.
5. David, H.W. and Mark, H.S. (2007). Climate and its Effect on Crop Production and Management, pp. 281-289.
6. Auta, S.J. and Dafwang, I.I. (2010). The agricultural development projects (ADPs) in Nigeria: Status and policy implications. *Journal of Agriculture and Biological Sciences*, vol. 6, no. 2, pp. 138-143.
7. Jifin, D.T. (2017). Impact of Climate Change on Agricultural Production by Farmers in Taraba State, Nigeria. *International Journal of Entrepreneurial Development, Education and Science Research*, vol. 4, no. 1, pp. 178-190.
8. Mahmood, H.U., Jerome, N. and Jane, J. (2014). Agricultural Management Strategy on Food Security in Taraba State. *Journal of Biology, Agriculture and Healthcare*, vol. 4, no. 8, pp. 29-31.
9. Angela, I.E. and Fidelis, H.B. (2013). Effects of Climate Change on Rice Farming in Ardo Kola Combination and Ordered Weighted Averaging. *Informatica*, vol. 33, no. 4, pp. 459-474.
10. Adebayo, A.A. and Umar, A.S. (1999). Hydrology and Water Resources in Adebayo, A. A. & Tukur, A. L. (Eds) Adamawa state in maps. *Geography Department, FUT, Yola & Paraclette Publishers, Yola*, pp. 17-19.
11. Taraba Youth Progressive Association of Nigeria (TYPA) (2009). Report on small scale environmental studies in Taraba state, pp. 11-19.
12. Emeka, D.O and Abbas, B. (2011). The geography of Taraba State, Nigeria. *Australia: LAP LAMBERT academic publishing LTD*, pp. 168-183.
13. Iloje, N.P. (2001). A new geography of Nigeria. *New Revised Edition. Longman Nigeria PLC*. 200p.

14. Adebayo, A.A. and Adebayo, E.F (1997). Forecasting the Yield of Upland Rice in Adamawa state, Nigeria on the basis of Agro-climatic Factors. *Malaysian Journal of tropical Geography*, vol. 28, no. 2, pp. 97-101.
15. Adebayo, A.A. (2000). Agro-climatic Classification of Adamawa State, Nigeria for Upland rice production. *The Nigerian Geographical Journal, New series*, vol. 3, no. 4, pp. 83-97.
16. Deotrefhy, K.D., Feroze, S.M., Ram, S. and Lala, I.P.R. (2017). Effect of Rainfall Variability on Rice Yield in North Eastern Hills of India. *Agricultural Research*, vol. 6, no. 4, pp. 341-346.
17. Mahmood, N., Ahmad, B., Hassan, S. and Bakhsh, K. (2012). Impact of temperature and precipitation on rice productivity in rice-wheat cropping system of Punjab Province. *The Journal of Animal & Plant Sciences*, vol. 22, no. 4, pp. 993-997.
18. Govinda, B. (2013). Effect of Precipitation and Temperature Variation On the Yield of Major Cereals in Dadeldhura District of Far Western Development Region, Nepal. *International Journal of Plant, Animal and Environmental Sciences*, vol. 3, no. 1, pp. 247-256.
19. Saliu, A.T., Jude, N.E., Taibat, M.Y., Alhassan, T.M. and Samuel, O.B. (2015). Rainfall Variability and Its Effect on Yield of Rice in Nigeria. *International Letters of Natural Sciences*, vol. 49, pp. 63-68.
20. Wahid, A., Gelani, S., Ashraf, M. and Foolad, M.R. (2007). Heat Tolerance in Plants: An overview. *Environmental and Experimental Botany*, vol. 61, pp. 199-223.
21. Farrell, T.C., Fox, K.M., Williams, R.L. and Fukai, S. (2006). Genotypic Variation for Cold Tolerance during Reproductive Development in Rice: Screening with cold air and Cold Water. *Field Crops Research*, vol. 98, pp. 178-194.
22. Yin, X., Kroff, M.J. and Goudriann, J. (1996). Differential Effects of Day and Night Temperature on Development to Flowering in Rice. *Annals of Botany*, vol. 77, pp. 203-213.
23. Sridevi, V. and Chellamuthu, V. (2015). Impact of weather on rice. *International Journal of Applied Research*, vol. 1, no. 9, pp. 825-831.
24. Expert System for Paddy-Water Requirement (2018). Received December, 4th 2018, from http://www.agritech.tnau.ac.in/expert_system/paddy/cultivationpractice.
25. Agropedia (2018). Irrigation water management in paddy. Received 4th December, 2018, from <http://agropedia.iitk.ac.in/content/irrigation-water-management-paddy>,
26. Jackson, M.B. (2013). The impact of flooding stress on plants and crops. URL Available at: [http://www.plantstress.com/Articles/\(waterlogging\)](http://www.plantstress.com/Articles/(waterlogging)).
27. Getachew, T.A. and Solomon, A.B. (2015). Land Suitability Analysis for Rice Production: A GIS Based Multi-Criteria Decision Approach. *American Journal of Geographic Information System*, vol. 4, no. 3, pp. 95-104.
28. Craufurd, P.Q. and Wheeler, T.R. (2009). Climate change and the flowering time of annual crops. *Journal of Experimental Botany*, vol. 60, pp. 2529-2539.
29. Ogbuene, E.B. (2010). Impact of Meteorological Parameters on Rice Yield; An Approach for Environmental Resources Sustainability in Ebonyi rice farmland, Nigeria. *Journal of Environmental Issues and Agriculture in Developmental Countries*, vol. 2, no. 3, pp. 103-116.
30. Arif, A., Harsh, B.P., Harish, K. and Surender, K. (2017). Effect of temperature and rainfall different on growth stages and yield of rice (*Oryza sativa* L.). *Journal of Pharmacognosy and Phytochemistry*, vol. 6, no. 1, pp. 331-332.
31. Mostafa M, Mohammad A. N and Mehdi M (2015). Assessing the Impact of Climate Variability on Rice Phenology. *Research Journal of Environmental Sciences*, vol. 9, pp. 296-301.

32. Tesfamariam, E.H., Annandale, J.G. and Steyn, J.M. (2010). Water stress effects on winter canola growth and yield. *Agron. Journal*, vol. 102, no. 2, pp. 658–666. <https://doi.org/10.2134/agronj2008.0043>.
33. Sawa, B. and Adebayo, A. (2011). The Impact of Climate on Precipitation Effectiveness Indices in Northern Nigeria. *Journal of Environmental and Earth Sciences*, pp. 481-486.
34. Moncumill, D.F. (2006). Climate Variability and Corn Yields in Semiarid Ceara; Brazil. *Journal of Applied Meteorology and Climatology*, vol. 46, pp.226 -280.